

## Accepted Manuscript

Electrochemical triggering of the Chardonnay wine metabolome

Chloé Roullier-Gall, Basem Kanawati, Daniel Hemmler, Gregory K. Druschel,  
Régis D. Gougeon, Philippe Schmitt-Kopplin


PII: S0308-8146(19)30224-9  
DOI: <https://doi.org/10.1016/j.foodchem.2019.01.149>  
Reference: FOCH 24244

To appear in: *Food Chemistry*

Received Date: 23 January 2018  
Revised Date: 16 January 2019  
Accepted Date: 22 January 2019

Please cite this article as: Roullier-Gall, C., Kanawati, B., Hemmler, D., Druschel, G.K., Gougeon, R.D., Schmitt-Kopplin, P., Electrochemical triggering of the Chardonnay wine metabolome, *Food Chemistry* (2019), doi: <https://doi.org/10.1016/j.foodchem.2019.01.149>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



## Electrochemical triggering of the Chardonnay wine metabolome

Chloé Roullier-Gall<sup>1,2,\*</sup>, Basem Kanawati<sup>2</sup>, Daniel Hemmler<sup>1,2</sup>, Gregory K. Druschel<sup>4</sup>, Régis D. Gougeon<sup>3</sup> and Philippe Schmitt-Kopplin<sup>1,2\*</sup>.

<sup>1</sup> Chair of Analytical Food Chemistry, Technische Universität München, Alte Akademie 10, 85354 Freising-Weihenstephan, Germany.

<sup>2</sup> Research Unit Analytical BioGeoChemistry, Department of Environmental Sciences, Helmholtz Zentrum München, Ingolstaedter Landstrasse 1, 85764 Neuherberg, Germany.

<sup>3</sup> Present adress : Univ. Bourgogne Franche-Comté, AgroSup Dijon, PAM UMR A 02.102, F-21000 Dijon, France. Institut Universitaire de la Vigne et du Vin Jules Guyot.

<sup>3</sup> Univ. Bourgogne Franche-Comté, AgroSup Dijon, PAM UMR A 02.102, F-21000 Dijon, France. Institut Universitaire de la Vigne et du Vin Jules Guyot.


<sup>4</sup> Department of Earth Sciences, Indiana University – Purdue University Indianapolis, IN 46202, United States.

### Corresponding authors:

\*Chloé Roullier-Gall: chloe.roullier-gall@u-bourgogne.fr

\*Philippe Schmitt-Kopplin: schmitt-kopplin@helmholtz-muenchen.de

**TOC graph:** The electrochemical oxidation of two bottle-aged wines, which only differ by the SO<sub>2</sub> dose added at pressing, was shown to induce different reactions related to this pre-fermentation winemaking process. Ultrahigh resolution mass spectrometry revealed the corresponding oxidation fingerprints and the impact of the SO<sub>2</sub> initially added to the must.



## Abstract

Oxidation of wine upon bottle ageing is a crucial matter of concern for the qualitative long-term storage of white wines. However, understanding the various molecular mechanisms potentially involved, which can impact the wine composition, requires that top-down analytical strategies are implemented. Here, we report the analysis of bottle aged Chardonnay wines made from the same must, but differing by the amount of SO<sub>2</sub> initially added to the must at pressing (0 and 8 g.hL<sup>-1</sup>). Metabolomics fingerprints obtained from electrochemical simulation of oxidative reactions were obtained by the coupling of either on-line or off-line electrochemical oxidation to FT-ICR-MS detection. We reveal that, whatever the electrochemical DC voltage, wines with initial SO<sub>2</sub> addition displayed molecular fingerprints, which remained more similar to the non-oxidized wine without initial SO<sub>2</sub> addition. We further show that a diversity of sulfur-containing compounds appeared to be the most sensitive to oxidation, whereas nitrogen-containing compounds were mostly formed.

**Keywords:** Chardonnay wine; oxidation; SO<sub>2</sub>; FT-ICR-MS; electrochemistry (EC).

## 1. Introduction

In contrast with many other food products, quality wines are designed to age for durations that can vary from weeks after bottling to potentially decades. If this intuitively applies for red wines, it is also the case for white wines, and in particular for well-known dry Chardonnay wines from Burgundy.

During its bottle ageing time, where various chemical changes occur (Hernanz et al., 2009; Kallithraka, Salacha, & Tzourou, 2009; Maury, Clark, & Scollary, 2010), the wine experiences three consecutive stages, corresponding to maturation and improved stability, organoleptic optimum, and decline (Arapitsas, Speri, Angeli, Perenzoni, & Mattivi, 2014). Through its antioxidant, antioxidasic and antiseptic properties, SO<sub>2</sub> is considered as an essential tool for promoting wine ageing, in particular for white wines, where it is used to prevent from browning (Li, Guo, & Wang, 2008), to slow down the decrease of desired aromas (Escudero, Asensio, Cacho, & Ferreira, 2002; Nikolantonaki, Magiatis, & Waterhouse, 2014; Roussis, Lambropoulos, & Tzimas, 2007), or more

generally to protect wine from autoxidation, which naturally occurs because of the intrinsic permeability to oxygen of stoppers (Karbowski, Mansfield, Barrera-García, & Chassagne, 2010; Ugliano, 2013; Waterhouse et al., 2016).

The antioxidant properties of SO<sub>2</sub> in wine have already been the subject of several studies, and if various mechanisms have been proposed, an integrated description of its impact on the chemistry of wine ageing remains to be elucidated. Results from experiments run on model wine have shown that oxygen is supposed to react only weakly with molecular SO<sub>2</sub>, but its reaction with hydrogen peroxide prevents ethanol oxidation via the Fenton reaction (Danilewicz, 2007; Waterhouse & Laurie, 2006). Due to its inherent nucleophilic character, the bisulfite ion (HSO<sub>3</sub><sup>-</sup>) easily reacts with carbonyl-containing compounds, in particular with quinones, either to form sulfonic adducts or to reduce them back to dihydroxybenzenes (Danilewicz, Seccombe, & Whelan, 2008; Maujean, 2001). However, it was also shown that these reactions depend on various parameters, including the structure of the involved phenolics, the concentration of transition element catalysts or the composition of organic acids (Danilewicz, 2011, 2014; Makhotkina & Kilmartin, 2013). Through untargeted LC-MS analyses, Arapitsas et al. showed that SO<sub>2</sub> added to wine takes part in various reactions including the formation of sulfonated adducts of indole-3-lactic hexoside, tryptophol, glutathione, cysteine and pantetheine (Arapitsas et al., 2016). They further demonstrated that the sulfonation of metabolites with an indole scaffold dominated in white wines (Arapitsas, Guella, & Mattivi, 2018). However, and except for indole-3-lactic acid, for which the adducts could reach concentrations up to 8 mg/L, most of these identified adducts appeared to exist at low concentrations in white wine, thus suggesting that the actual chemistry related to SO<sub>2</sub> combinations certainly involves a significantly higher diversity of wine metabolites.

SO<sub>2</sub> can be added at different times of the winemaking of whites, starting with the protection of the must before, during, or immediately after pressing. Then if malolactic fermentation is desired, SO<sub>2</sub> is added at the end to protect the wine during ageing, which can proceed in barrels or in tanks. Finally, SO<sub>2</sub> is added before bottling. Although the first must addition assumes mostly antioxidasic and antiseptic purposes, several studies have shown that it has an impact on the long-term oxidative

stability of white wines (Cheynier, Masson, Rigaud, & Moutounet, 1993; Ugliano, 2013)(Cejudo-Bastante, Hermosín-Gutiérrez, Castro-Vázquez, & Pérez-Coello, 2011; Schneider, 1998) (Cheynier, Souquet, Samson, & Moutounet, 1991). Recently, and through untargeted Fourier transform ion cyclotron resonance mass spectrometry (FT-ICR-MS) metabolomics, we have further shown that hundreds of masses and likely thousands of must and wine compounds are actually involved in chemical mechanisms related to SO<sub>2</sub> addition to the must, including amino acids, organic acids, carbohydrates or polyphenols, and that these compounds, which are still present after several years of bottle ageing, allow discrimination of bottle aged white wines according to the SO<sub>2</sub> concentration added at pressing (Chloé Roullier-Gall et al., 2017).

However, in a context where there is a trend over recent years to limit the use of SO<sub>2</sub> (Ancín-Azpilicueta, Jiménez-Moreno, Moler, Nieto-Rojo, & Urmeneta, 2016), and consequently an increasing demand for alternatives, understanding molecular mechanisms by which SO<sub>2</sub> operates as antioxidant during the long-term bottle ageing of wine is a prerequisite. Measuring an antioxidant capacity of wine and relating it to the many potentially involved antioxidant compounds remains a challenge. In order to get antioxidant capacity of wine, electrochemical methods have proved to be interesting alternative strategies to classical spectrophotometric methods, which monitor the formation or the disappearance of radical-related chromogenic compounds (Hoyos-Arbeláez, Vázquez, & Contreras-Calderón, 2017). Different authors have indeed attempted to provide estimated antioxidant capacity index or electrochemical index of wines, based in particular on voltammetric analyses (Gao et al., 2015; Kilmartin, Zou, & Waterhouse, 2001; Lino et al., 2014). It has been shown in particular, that anodic currents related to the oxidation of selected polyphenols could be selectively monitored (Makhotkina & Kilmartin, 2009). Considering representative wine polyphenols (catechin, caffeic acid and quercetin) in model wine conditions, these authors further showed that in the presence of SO<sub>2</sub>, electrochemical oxidation could lead to semiquinone radicals, which can be either reduced back to the original polyphenol, or further oxidized to the quinone form. However, in combination with HPLC-MS analyses, they demonstrated that depending on the polyphenol structure, quinones could either react with SO<sub>2</sub> and produce new derivatives, or undergo chemical reactions to produce several new

compounds without SO<sub>2</sub> combination (Makhotkina & Kilmartin, 2013). This is one of very few examples combining electrochemical and mass spectrometric analyses of wine related samples; it also points to the limits of targeted molecular analyses, which are far from characterizing the more complete diversity of wine compounds potentially involved in the antioxidant process.

In this paper, combination of electrochemical oxidation strategies and ultra-high resolution FT-ICR-MS were used to characterize the antioxidant property of white wines from an untargeted molecular point of view. Seven year old bottled Chardonnay white wines, only differing by the concentration of SO<sub>2</sub> added to the must at pressing, were considered here in order to account for the intrinsic chemical diversity related to the various chemical pathways, through which SO<sub>2</sub> is involved during the first years of bottle ageing.

## **2. Experimental**

### **2.1 Wine Samples.**

Chardonnay wines, from Montagny appellation in Burgundy, France and from the 2007 vintages, were analyzed. For each of the conditions, 3 biological replicates (bottles) were analyzed. These wines were originated from one single must, which was treated with two different concentrations of sulfur dioxide (0 and 8 g.hL<sup>-1</sup>) immediately after pressing, and before alcoholic fermentation. All wines were bottled after supplementation of a constant SO<sub>2</sub> addition to ensure a free SO<sub>2</sub> concentration of the order of 35 mg/L. Free and total SO<sub>2</sub> concentrations were measured after 6 months of bottle ageing. Wines with 0 g.hL<sup>-1</sup> SO<sub>2</sub> addition at pressing contained in average 17.5 mg/L free SO<sub>2</sub> and 91 mg/L total SO<sub>2</sub> and wine with 8 .hL<sup>-1</sup> SO<sub>2</sub> addition at pressing contained 17.5 mg/L free SO<sub>2</sub> and 120.5 mg/L total SO<sub>2</sub>. Electro oxidation experiments were done between January and June 2015.

### **2.2 Oxidation by Electrochemistry.**

**2.2.1 On-line electrochemistry:** An electrochemical  $\mu$ -PrepCell™ Roxy (Antec, Netherlands)

equipped with a conductive diamond (MD) working electrode was used (Supplemental Figure 1). A Pd/H<sub>2</sub> electrode was used as a reference electrode (RE) and titanium electrode as a counter electrode. Samples were analyzed at three different electrical DC potentials (400 mV, 800 mV and 1200 mV). Additionally the samples were mass spectrometrically acquired without applying a potential (0 mV). The liquid outlet of the electrochemical cell was connected directly to the electrospray source of a FT-ICR-MS (Bruker Daltonics, Germany), which is used to record mass spectra. Diluted Wine (1/20) in a solvent mixture of (50:50 water : methanol) was pumped through the electrochemical cell at a flow rate of 10  $\mu\text{L}\cdot\text{min}^{-1}$ . The cell volume is 11  $\mu\text{L}$  and the working electrode area is 1.9  $\text{cm}^2$ . The residence time of the wine in the  $\mu$ -PrepCell™ Roxy cell was 1 min and 6 s.

**2.2.2 Off-line electrochemistry:** EC experiments were acquired using a DLK-70 Web-Pstat™

Potentiostat (Analytical Instrument Systems, Flemington, USA) (Supplemental Figure 1). A self-made copper electrode was used as working electrode (WE), an Ag/AgCl electrode was used as a reference electrode (RE) and Pt electrode as a counter electrode. The working electrode area is 1.3  $\text{cm}^2$ . The surface area is calculated by assuming cylindrical geometry of wires measured to be 75  $\mu\text{m}$  in diameter each. For each sample, a new copper WE electrode was used in order to ensure same conditions and unmodified surface. 2 ml of each sample were treated with constant potentials (500 mV, 1000 mV and 1500 mV) applied between the copper WE and Ag/AgCl RE. Wine samples were collected after 10 min of EC treatment. After electrochemical treatment the samples were analyzed by direct infusion FT-ICR-MS.

### **2.3 Ion Cyclotron Resonance Fourier Transform Mass Spectrometer.**

Ultrahigh resolution mass spectra were acquired using a solariX FT-ICR-MS instrument (Bruker Daltonik, Bremen, Germany) equipped with a 12 Tesla superconducting magnet and an Apollo II electrospray ionization source operated in the negative ionization mode. 50  $\mu\text{L}$  of the samples were diluted in 1 ml of methanol prior to injection and introduced into the micro electrospray source at a

flow rate of  $120\mu\text{L}\cdot\text{h}^{-1}$  using a syringe pump. The MS was externally calibrated based on cluster ions of arginine (10 ppm). The mass range is  $m/z$  100 to 1000 Da. 300 scans were accumulated in an acquisition for each sample and spectra were acquired with a time domain of 4 mega-words. Spectra were internally recalibrated using a composed list of fatty acids and recurrent wine compounds, applying linear calibration until  $m/z$  1000, with mass errors below 50 ppb. Peaks with a signal to noise ratio (S/N) of 4 and higher were used for further data processing. In addition to an automated theoretical isotope pattern comparison, the generated formulas were further validated by setting sensible chemical constraints (N rule; O/C ratio  $\leq 1$ ; H/C ratio  $\leq 2n+2$ ; element counts:  $C \leq 100$ ,  $H \leq 200$ ,  $O \leq 80$ ,  $N \leq 3$ ,  $S \leq 3$  and  $P \leq 1$ ). FT-ICR-MS peak alignment and filtering of masses were performed in MS Excel 2010 (Microsoft, Redmond, USA) with maximum error thresholds of 1 ppm and filtered for masses occurring at least in 10% of all samples (Roullier-Gall, Witting, Gougeon, & Schmitt-Kopplin, 2014).

### 3. Results and discussion

#### 3.1 On-line EC-MS simulation of oxidative ageing

Initially, on-line EC-MS experiments were performed to obtain information on the electrochemical oxidation behavior of white wine depending on the  $\text{SO}_2$  dose. Two wines, from identical musts, treated immediately after pressing with two different concentrations of sulfur dioxide (0 and  $8\text{ g}\cdot\text{hL}^{-1}$ ) were subjected to DC potentials (from 0 to 1200 mV). A commercial electrochemical flow-through cell ( $\mu\text{-PrepCell}^{\text{TM}}$ ) including a conductive diamond working electrode, a palladium counter electrode and a Pd/ $\text{H}_2$  reference electrode was used. The hierarchical cluster analysis based on the matrix of all masses for wines with 0 and  $8\text{ g}\cdot\text{hL}^{-1}$   $\text{SO}_2$  initially added to the must is presented in Figure 1. The  $\text{SO}_2$  dose (0 or  $8\text{ g}\cdot\text{hL}^{-1}$   $\text{SO}_2$ ) was used as initial condition prior to apply any electrical potential. Samples were clustered into two groups (Figure 1A). The first group corresponds to sulfur treated samples ( $8\text{ g}\cdot\text{hL}^{-1}$   $\text{SO}_2$  in blue), independently of the applied voltage (0, 400, 800 and 1200 mV) in addition to the zero  $\text{SO}_2$  added sample ( $0\text{ g}\cdot\text{hL}^{-1}$  in grey) prior electrochemical oxidation (0 mV). The second group

corresponds to wines without SO<sub>2</sub> addition at pressing (0 g.hL<sup>-1</sup> in grey) after electrochemical oxidation (400, 800 and 1200 mV) (Figure 1A).

Wines without SO<sub>2</sub> addition at pressing showed significant compositional changes (chemical reactivity) when a DC potential of 400 mV is applied. This was the minimum required DC voltage, below which no significant modifications could be observed with our FT-ICR-MS detection, although some phenolics can oxidize already below that value (Makhotkina & Kilmartin, 2013; Ugliano, 2016). With increasing DC voltages from 400mV to 1200mV, more compositional changes and sharper signal intensity changes could be experimentally observed with the on-line EC system (Figure and Supplemental Figure 2). Wines without SO<sub>2</sub> addition at pressing (0 g.hL<sup>-1</sup> of SO<sub>2</sub> in grey) showed the highest internal variation of elemental chemical compositions due to electrical potential (Figure 1A). The composition of sulfur treated samples (8 g.hL<sup>-1</sup> of SO<sub>2</sub> in blue) appeared to be less impacted by electrochemical oxidation (Figure 1A). Additional conclusions can be elucidated considering the masses that show a significant change in peak intensity after electrochemical treatment (van Krevelen diagrams shown in Figure 1B and C).

Some metabolites were characterized by decreasing intensities as a result of the oxidation process and some others were newly formed (Figure 1B and C, supplemental Figure 3). Masses, whose signal intensity decreased as a function of the applied electrical potential, were considered reactive compounds toward electro-oxidation. These reactive compounds, which are located in the polyphenol and peptide regions, as seen in the Van Krevelen diagrams, represent mostly sulfur containing species (CHOS and CHNOS) (Figure 1 and Supplementary Table 1). Masses, whose signal intensities significantly increase as a function of applied electrochemical DC potential, are considered electro-oxidation products. These products are mainly nitrogen containing compounds (CHON, orange) and CHO compounds (CHO in blue) (Figure 1 and Supplementary Table 1). According to Supplementary Table 1, a larger number of reactive compounds were found in wines without SO<sub>2</sub> addition at pressing (385 decreasing masses) compared to wines for which the must received 8 g.hL<sup>-1</sup> of SO<sub>2</sub> (222 decreasing masses). Interestingly, the number of oxidation products was of the same order for the two

types of wines (171 increasing masses for wine without SO<sub>2</sub> addition at pressing and 157 for SO<sub>2</sub> treated musts) (Supplementary Table 1). The comparison of van Krevelen diagrams of reactive compounds and products toward electro-oxidation highlighted a shift from low to higher O/C ratio as expected for oxidation processes (Roullier-Gall et al., 2016). Specific formulas of reactive compounds are distributed along O/C ratio from 0.2 to 1 whereas those of products are mainly located between 0.4 and 1 (Figure 1B and C). Altogether, bearing in mind that all wines received SO<sub>2</sub> at bottling, and that high additions of SO<sub>2</sub> to the must tend to produce wines with higher levels of combined SO<sub>2</sub> (as shown by SO<sub>2</sub> levels after bottling), our results suggest first that those compounds reacting to electrochemical oxidation are not necessarily those, which had reacted with SO<sub>2</sub> added to the must, and second that those compounds reacting to electrochemical oxidation are in particular sulfur- and nitrogen-containing compounds, which may either be native compounds from the grape or produced by fermentation. Therefore, these results provide unprecedented insights into the distinct chemical mechanisms involved in the reactivity toward electrochemical oxidation of bottle aged Chardonnay white wines, as a consequence of distinct levels of SO<sub>2</sub> initially added to the must.

Combining the on-line EC device with FT-ICR-MS is a fast and simple issue for getting a quick insight into the oxidative resistance of wines. On the other hand, the off-line system allows for a better control of the experimental setup. For example, the on-line system cannot tolerate high concentration of infused wine due to oversaturation of the working electrode and also because the electrospray ionization source, which is directly connected to the on-line system, cannot deal with high ion abundance which would consequently cause ICR cell ion oversaturation problems. For the off-line electrochemical approach, it is possible to use larger wine concentrations and flexibly extended oxidation periods for achieving better oxidation efficiency. The final oxidation products can be consequently further diluted in methanol to suit the electrospray source, and ICR-MS operation requirements. The extent of electrochemical oxidation is easily controllable by manipulating the time of electrochemical treatment and the ratio of electroactive surface area to fluid (i.e. higher surface area yields a faster reaction rate). For example, the time constraint in the on-line system is 1.1 min, whereas there is actually no time constraint for the off-line EC approach.

### 3.2 Off-line versus on-line coupling

We compared the on-line system and off-line system combined with FT-ICR-MS at 0 and 1000 mV for the same wine samples with and without SO<sub>2</sub> addition at pressing (Supplementary Table 2). With increasing DC voltages from 400 mV to 1200 mV, more compositional changes and sharper signal intensity changes had already been experimentally observed with the on-line system (Supplementary Figure 3). Based on these results, we used a standard potential of 1000 mV for off-line EC oxidation experiments. Using 10 min off-line electrochemical oxidation led to much more oxidation products with high m/z signal intensities (Figure 2). For samples without SO<sub>2</sub> addition (0 g.hL<sup>-1</sup>), and using the off-line system, the total number of detected masses was higher, 7,637 versus 5,710 formulas for off-line versus on-line, independently of the DC voltage applied (Supplementary Table 2 and Figure 2). Histograms of unique formulae, colored according to chemical families (CHO, CHOS, CHNO and CHNOS), showed differences in the number of compounds for “cell off” (0 mV) and “cell on” (DC potential of 1000mV) as a function of the SO<sub>2</sub> dose (Figure 2). New compounds have been formed during the simulated oxidation (Figure 2 and Supplementary Table 2). Venn diagrams of the wine composition for “cell off” and “cell on” conditions confirm that numerous compounds appear and disappear during the simulated oxidation. According to venn diagrams and histograms of elemental formulas sorted by chemical families, similarities could be observed between both systems. In agreement with the literature (Arapitsas et al., 2018; Ugliano, 2016), masses corresponding to catechin/epicatechin and caffeic acid are examples for reactive compounds for both on-line and off-line systems, but with distinct behaviours according to the SO<sub>2</sub> initially added to the must (Supplementary Table 3). The concentration decrease for these two compounds appeared to be more significant for wines which had no SO<sub>2</sub> addition to the must (Supplementary Figure 4), consistently with the fact that these reactive compounds are less concentrated in these wines, and/or that some new compounds formed upon initial addition of SO<sub>2</sub> to the must exhibit higher antioxidant efficiency than these polyphenols. On the basis that a 1000 mV voltage would likely promote the oxidation of the A ring of catechin/epicatechin, such results shows that adding SO<sub>2</sub> to the must would limitate this reaction both for the monomer and the adduct. A close number of elemental formulas impacted by DC

voltage are detected using on-line and off-line systems and sulfur containing compounds (CHOS) were mostly consumed, whereas nitrogen containing compounds (CHON and CHONS) were mostly created regardless of the oxidation system used (Figure 2).

### 3.3 Resistance to electrochemical oxidation chemical fingerprints

Simulated electrochemical oxidation using the Off-line system coupled to FT-ICR-MS, did not only induce a complete appearance or disappearance of some compounds. It also changed relative concentrations of many others, represented by changes in peak intensity. More than 60% of the total masses were present in wine samples regardless of experimental conditions (cell off and on). As example, for wines without SO<sub>2</sub> added, 6662 masses, out of 7637, were present in both conditions (cell off and on) (Figure 2). However, changes in intensity induced by an electrochemical potential could be observed also for these omnipresent compounds. Specific masses with significantly higher intensities for “cell off” and “cell on” (1000 mV) samples were extracted for wines without SO<sub>2</sub> added (Figure 3). The “cell off” samples were discriminated based on 646 masses with significantly higher intensities than for “cell on” samples (ANOVA test, p-values  $\leq 0.01$ ). The corresponding elemental formulae were represented in van Krevelen diagrams according to H/C and O/C ratio and histograms according to chemical family (CHO, CHOS, CHNO and CHNOS) (Figure 3). Discriminant masses for “cell off” showed a large variety of elemental formulas mainly with sulfur and nitrogen containing compounds (in red) consistently with previous results (Chloé Roullier-Gall et al., 2017). “Cell on” (1000 mV) samples were characterized by 1120 masses, which were significantly higher in signal intensities when compared to the wine samples prior to electro-oxidation (cell off) (ANOVA test, p-values  $\leq 0.01$ ). A large number of sulfur containing compounds (CHOS and CHONS) showed variation of intensity after the electrochemical oxidation and were found in areas corresponding to peptides, amino acids, and Maillard reaction products (Hemmler et al., 2017), suggesting the onset of a diversity of chemical reactions, which may include for instance Strecker degradation or Michael addition of amino acids/peptides to quinones (Bittner, 2006), when

polyphenols are consumed (Supplementary Table 3). In agreement with on-line experiments above (Figure 2), discriminant masses for oxidized samples (cell on in Figure 3B) were characterized by a higher number of sulfur containing compounds (46 CHOS compounds, in green) and nitrogen and sulfur containing compounds (133 CHNOS compounds, in red). Most of these masses were concentrated in the area of carbohydrates, and phenolic compounds, suggesting a possible sulfonation of these compounds. The comparison of these results to a recent study on the natural oxygenation of champagne wines (Roullier-Gall et al., 2016) revealed 38 common markers between low oxygenated champagne wines and markers of samples prior to electro-oxidation (cell off markers) and 13 common markers between high oxygenated champagne wines and markers of samples after electro-oxidation (Supplementary Table 4).

During electro oxidation there is also a noticeable hydrogen loss. A decreasing number of hydrogen atoms in a molecule increases unsaturation and hence leads to higher double bond equivalent (DBE) values. However, since large molecules can potentially contain more double bonds and rings than small molecules, in this study the DBE was normalized to the total number of carbon atoms (DBE/C). Thus, the DBE/C ratio will increase with decreasing H/C ratio (Koch & Dittmar, 2006). Based on the DBE/C ratio comparison between “cell on” and “cell off” markers, structural information can be extracted. Indeed “cell off” markers appeared more confined in the small mass range between  $m/z$  100-500 amu and are characterized by low DBE/C ratio, while “cell on” markers were more abundant in higher masses ( $m/z$  250-550 amu) and characterized by higher DBE/C values (Figure 3C). “Cell off” markers would be described as non-oxidized compounds whereas “cell on” markers would therefore correspond to oxidized compounds, as confirmed by the wine oxidation mechanism leading to the presence of higher masses with a higher number of double bonds (Bittner, 2006; Nikolantonaki et al., 2014; Waterhouse & Laurie, 2006; Wildenradt & Singleton, 1974). A further oxidation signature was provided by the number of detected O-containing formulas as a function of  $m/z$  for “cell on” and “cell off” markers, which is clearly higher for the former (Figure 3D, Supplementary Figure 5 and 6 and supplementary table 5). Interestingly, a count of the number of possible transformations connecting the detected masses, and corresponding for instance to hydroxylation,

hydro-Peroxidation, dehydrogenation, hydrolysis/condensation, thiolation and direct sulfonation showed for instance, that (de)hydroxylation ( $\pm\text{O}$ ) transformations would indeed be slightly more observed after electro oxidation (cell on) of wines without  $\text{SO}_2$  added to the must, and even more observed after electro oxidation of wines with  $8 \text{ g.hL}^{-1} \text{SO}_2$  added to the must (Supplementary table 5). However, thiolation ( $\pm\text{S}$ ) and sulfonation ( $\pm\text{SO}_3$ ) transformations would exhibit a contrasted behavior, where they are similarly and significantly more observed after electro oxidation (cell on) of wines without  $\text{SO}_2$  added to the must, but slightly less observed after electro oxidation of wines with  $8 \text{ g.hL}^{-1} \text{SO}_2$  added to the must. In contrast, sulfonation ( $\pm\text{H}_2\text{SO}_3$ , corresponding to the addition of sulfites to carbonyls) would exhibit the opposite trend with much more transformations observed for cell-on (Supplementary Table 5). The reactivity of sulfonated compounds is indeed central to this study and these results illustrate how the resistance of wines to electro oxidation involves distinct reactional mechanisms depending on the fact that musts were initially protected with  $\text{SO}_2$  or not. A closer look at masses corresponding to glutathione and glutamylcystein, and their sulfonated adducts further showed that such antioxidants exhibit distinct consumption behaviors upon electro oxidation (Supplementary Figures 5, 6 and 7). If all of these compounds appeared to be consumed upon electro oxidation (cell on), the consumption of glutathione and glutamylcystein was increased for wines with  $8 \text{ g.hL}^{-1} \text{SO}_2$  added to the must, whereas it is the opposite when their sulfonated adducts are considered (Supplementary Figure 7), *i.e.* sulfonated adducts formed upon  $\text{SO}_2$  addition at bottling only ( $0 \text{ g.hL}^{-1} \text{SO}_2$  added to the must) would be relatively more consumed after electro oxidation, than sulfonated adducts formed upon  $\text{SO}_2$  addition both at pressing and bottling ( $8 \text{ g.hL}^{-1} \text{SO}_2$  added to the must). Our results thus shows that such thiol S-sulfonate adducts formed upon ageing (Arapitsas et al., 2016; Bekker, Kreitman, Jeffery, & Danilewicz, 2018) may contribute to the overall antioxidant resistance of wines.

It must be noted though, that wines studied here were bottle aged for 6 years, and consequently may have already gone through a natural slow oxidation process, which means that cell off markers may already exhibit oxidized traits. The richness of information, which could be deduced for the decreasing and increasing number of annotated detected compounds, which are distributed along

several regions in the shown Van Krevelen diagram (Figure 3) confirmed that oxidation not only impacts polyphenolic compounds but also modify amino acids, carbohydrates and sulfur containing compounds.

#### 4. Conclusions

In this work, the electrochemical simulation of oxidative reactions in wine coupled to FT-ICR-MS detection was investigated in order to unravel the diversity and nature of metabolites involved in the resistance of bottled wine towards electrochemical oxidation. Through the comparison of bottle aged chardonnay wines, which only differed by the amount of SO<sub>2</sub> added to the must at pressing, this study addressed for the first time the impact of SO<sub>2</sub> addition to the must, from a bottled wine ageing chemistry point of view. The on-line coupling of electrochemical oxidation and FT-ICR-MS detection gives a rapid access to wine electrochemical oxidation fingerprints but with low oxidation efficiency if coupled to high-resolution MS, whereas an off-line EC system allows for a more detailed investigation of oxidation products, due to time requirements which are disconnected for oxidation and MS acquisition. Besides known metabolites such as catechin/epicatechin and caffeic acid, sulfur-containing compounds (CHOS) appeared to be the most reactive (concentration decreasing with electrochemical oxidation), whereas nitrogen-containing compounds were mostly formed. From a methodological point of view, these results reveal an accessible chemodiversity associated with electrochemical oxidation. As such, they constitute an unprecedented basis for top down exploration of potential anti oxidation biomarkers and subsequent antioxidant alternatives to SO<sub>2</sub>.

#### Acknowledgment

We are very grateful to the wine producer, Domaine Sauvestre, who provided the samples, and to the Bureau Interprofessionnel des Vins de Bourgogne (BIVB) for the experimental set up. We warmly thank Antec Scientific for the lending of the ROXY EC system and Dr. Daniel Vetter for his help.

## References

- Ancín-Azpilicueta, C., Jiménez-Moreno, N., Moler, J. A., Nieto-Rojo, R., & Urmeneta, H. (2016). Effects of reduced levels of sulfite in wine production using mixtures with lysozyme and dimethyl dicarbonate on levels of volatile and biogenic amines. *Food Additives & Contaminants: Part A*, 33(10), 1518–1526.
- Arapitsas, P., Guella, G., & Mattivi, F. (2018). The impact of SO<sub>2</sub> on wine flavanols and indoles in relation to wine style and age. *Scientific Reports*, 8(1), 858. <https://doi.org/10.1038/s41598-018-19185-5>
- Arapitsas, P., Speri, G., Angeli, A., Perenzoni, D., & Mattivi, F. (2014). The influence of storage on the “chemical age” of red wines. *Metabolomics*, 1–17. <https://doi.org/10.1007/s11306-014-0638-x>
- Arapitsas, P., Ugliano, M., Perenzoni, D., Angeli, A., Pangrazzi, P., & Mattivi, F. (2016). Wine metabolomics reveals new sulfonated products in bottled white wines, promoted by small amounts of oxygen. *Journal of Chromatography A*, 1429, 155–165. <https://doi.org/10.1016/j.chroma.2015.12.010>
- Bekker, M. Z., Kreitman, G. Y., Jeffery, D. W., & Danilewicz, J. C. (2018). Liberation of Hydrogen Sulfide from Dicysteinyl Polysulfanes in Model Wine. *Journal of Agricultural and Food Chemistry*, 66(51), 13483–13491. <https://doi.org/10.1021/acs.jafc.8b04690>
- Bittner, S. (2006). When quinones meet amino acids: chemical, physical and biological consequences. *Amino Acids*, 30(3), 205–224. <https://doi.org/10.1007/s00726-005-0298-2>
- Cejudo-Bastante, M. J., Hermosín-Gutiérrez, I., Castro-Vázquez, L. I., & Pérez-Coello, M. S. (2011). Hydroxylation and Bottle Storage of Chardonnay White Wines: Effects on Color-Related Phenolics, Volatile Composition, and Sensory Characteristics. *Journal of Agricultural and Food Chemistry*, 59(8), 4171–4182. <https://doi.org/10.1021/jf104744q>
- Cheyrier, V., Masson, G., Rigaud, J., & Moutounet, M. (1993). Estimation of must oxidation during pressing in Champagne. *American Journal of Enology and Viticulture*, 44(4), 393–399.
- Cheyrier, V., Souquet, J.-M., Samson, A., & Moutounet, M. (1991). Hydroxylation: influence of various oxygen supply levels on oxidation kinetics of phenolic compounds and wine quality. *Vitis*, 30(2), 107–115.
- Danilewicz, J. C. (2007). Interaction of Sulfur Dioxide, Polyphenols, and Oxygen in a Wine-Model System: Central Role of Iron and Copper. *American Journal of Enology and Viticulture*, 58(1), 53–60.
- Danilewicz, J. C. (2011). Mechanism of autoxidation of polyphenols and participation of sulfite in wine: key role of iron. *American Journal of Enology and Viticulture*, ajev–2011.
- Danilewicz, J. C. (2014). Role of tartaric and malic acids in wine oxidation. *Journal of Agricultural and Food Chemistry*. <https://doi.org/10.1021/jf5007402>
- Danilewicz, J. C., Secombe, J. T., & Whelan, J. (2008). Mechanism of interaction of polyphenols, oxygen, and sulfur dioxide in model wine and wine. *American Journal of Enology and Viticulture*, 59(2), 128–136.
- Escudero, A., Asensio, E., Cacho, J., & Ferreira, V. (2002). Sensory and chemical changes of young white wines stored under oxygen. An assessment of the role played by aldehydes and some other important odorants. *Food Chemistry*, 77(3), 325–331.

Gao, F., Zheng, D., Tanaka, H., Zhan, F., Yuan, X., Gao, F., & Wang, Q. (2015). An electrochemical sensor for gallic acid based on Fe<sub>2</sub>O<sub>3</sub>/electro-reduced graphene oxide composite: Estimation for the antioxidant capacity index of wines. *Materials Science and Engineering: C*, *57*, 279–287.

Hemmler, D., Roullier-Gall, C., Marshall, J. W., Rychlik, M., Taylor, A. J., & Schmitt-Kopplin, P. (2017). Evolution of Complex Maillard Chemical Reactions, Resolved in Time. *Scientific Reports*, *7*.  
<https://doi.org/10.1038/s41598-017-03691-z>

Hernanz, D., Gallo, V., Recamales, Á. F., Meléndez-Martínez, A. J., González-Miret, M. L., & Heredia, F. J. (2009). Effect of storage on the phenolic content, volatile composition and colour of white wines from the varieties Zalema and Colombar. *Food Chemistry*, *113*(2), 530–537.

Hoyos-Arbeláez, J., Vázquez, M., & Contreras-Calderón, J. (2017). Electrochemical methods as a tool for determining the antioxidant capacity of food and beverages: A review. *Food Chemistry*, *221*, 1371–1381.

Kallithraka, S., Salacha, M. I., & Tzourou, I. (2009). Changes in phenolic composition and antioxidant activity of white wine during bottle storage: Accelerated browning test versus bottle storage. *Food Chemistry*, *113*(2), 500–505.

Karbowiak, T., Mansfield, A. K., Barrera-García, V. D., & Chassagne, D. (2010). Sorption and diffusion properties of volatile phenols into cork. *Food Chemistry*, *122*(4), 1089–1094.  
<https://doi.org/10.1016/j.foodchem.2010.03.089>

Kilmartin, P. A., Zou, H., & Waterhouse, A. L. (2001). A Cyclic Voltammetry Method Suitable for Characterizing Antioxidant Properties of Wine and Wine Phenolics. *Journal of Agricultural and Food Chemistry*, *49*(4), 1957–1965. <https://doi.org/10.1021/jf001044u>

Koch, B. P., & Dittmar, T. (2006). From mass to structure: an aromaticity index for high-resolution mass data of natural organic matter. *Rapid Communications in Mass Spectrometry*, *20*(5), 926–932.

Li, H., Guo, A., & Wang, H. (2008). Mechanisms of oxidative browning of wine. *Food Chemistry*, *108*(1), 1–13.

Lino, F. M. A., de Sá, L. Z., Torres, I. M. S., Rocha, M. L., Dinis, T. C. P., Ghedini, P. C., ... Gil, E. S. (2014). Voltammetric and spectrometric determination of antioxidant capacity of selected wines. *Electrochimica Acta*, *128*, 25–31.

Makhotkina, O., & Kilmartin, P. A. (2009). Uncovering the influence of antioxidants on polyphenol oxidation in wines using an electrochemical method: Cyclic voltammetry. *Journal of Electroanalytical Chemistry*, *633*(1), 165–174.

Makhotkina, O., & Kilmartin, P. A. (2013). Electrochemical Oxidation of Wine Polyphenols in the Presence of Sulfur Dioxide. *Journal of Agricultural and Food Chemistry*, *61*(23), 5573–5581.  
<https://doi.org/10.1021/jf400282z>

Maujean, A. (2001). The chemistry of sulphur in musts and wines. *JOURNAL INTERNATIONAL DES SCIENCES DE LA VIGNE ET DU VIN*, *35*(4), 171–194.

Maury, C., Clark, A. C., & Scollary, G. R. (2010). Determination of the impact of bottle colour and phenolic concentration on pigment development in white wine stored under external conditions. *Analytica Chimica Acta*, *660*(1), 81–86.

Nikolantonaki, M., Magiatis, P., & Waterhouse, A. L. (2014). Measuring protection of aromatic wine thiols from oxidation by competitive reactions vs wine preservatives with ortho-quinones. *Food Chemistry*, *163*, 61–67. <https://doi.org/10.1016/j.foodchem.2014.04.079>

- Roullier-Gall, C., Witting, M., Gougeon, R. D., & Schmitt-Kopplin, P. (2014). High precision mass measurements for wine metabolomics. *Frontiers in Chemistry*, 2, 102. <https://doi.org/10.3389/fchem.2014.00102>
- Roullier-Gall, C., Witting, M., Moritz, F., Gil, R. B., Goffette, D., Valade, M., ... Gougeon, R. D. (2016). Natural oxygenation of Champagne wine during ageing on lees: A metabolomics picture of hormesis. *Food Chemistry*, 203, 207–215. <https://doi.org/10.1016/j.foodchem.2016.02.043>
- Roullier-Gall, Chloé, Hemmler, D., Gonsior, M., Li, Y., Nikolantonaki, M., Aron, A., ... Schmitt-Kopplin, P. (2017). Sulfites and the wine metabolome. *Food Chemistry*, 237, 106–113.
- Roussis, I. G., Lambropoulos, I., & Tzimas, P. (2007). Protection of Volatiles in a Wine with Low Sulfur Dioxide by Caffeic Acid or Glutathione. *American Journal of Enology and Viticulture*, 58(2), 274–278.
- Schneider, V. (1998). Must Hyperoxidation: A Review. *American Journal of Enology and Viticulture*, 49(1), 65–73.
- Ugliano, M. (2013). Oxygen Contribution to Wine Aroma Evolution during Bottle Aging. *Journal of Agricultural and Food Chemistry*, 61(26), 6125–6136. <https://doi.org/10.1021/jf400810v>
- Ugliano, M. (2016). Rapid fingerprinting of white wine oxidizable fraction and classification of white wines using disposable screen printed sensors and derivative voltammetry. *Food Chemistry*, 212, 837–843. <https://doi.org/10.1016/j.foodchem.2016.05.156>
- Waterhouse, A. L., Frost, S., Ugliano, M., Cantu, A. R., Currie, B. L., Anderson, M., ... others. (2016). Sulfur Dioxide–Oxygen Consumption Ratio Reveals Differences in Bottled Wine Oxidation. *American Journal of Enology and Viticulture*, ajev–2016.
- Waterhouse, A. L., & Laurie, V. F. (2006). Oxidation of Wine Phenolics: A Critical Evaluation and Hypotheses. *American Journal of Enology and Viticulture*, 57(3), 306–313.
- Wildenradt, H. L., & Singleton, V. L. (1974). The production of aldehydes as a result of oxidation of polyphenolic compounds and its relation to wine aging. *American Journal of Enology and Viticulture*, 25(2), 119–126.

**Figure 1:** Evolution of the composition of Chardonnay wines after electrochemical oxidation treatment. (A) Hierarchical cluster analysis of wine without SO<sub>2</sub> addition at pressing and wines treated with 8 g.hL<sup>-1</sup> SO<sub>2</sub> used as initial condition prior to applied electrical potential (0, 400, 800 and 1200 mV) using On-line system. Van Krevelen diagrams and histograms showing metabolites that have been strongly affected by electrical potential for (B) SO<sub>2</sub> treated samples and (C) samples without SO<sub>2</sub> addition at pressing. Van Krevelen diagrams from left show metabolites with decreasing signal intensities and from right show newly appeared metabolites as a result of oxidation. Elemental formulas in the van Krevelen diagram are colored according to their composition and are sized according to their intensity, CHO in blue, CHOS in green, CHON in orange and CHONS in red.

**Figure 2:** Composition of wines treated with on-line and off-line system, both with FT-ICR-MS detection. Venn diagrams of the total white wine composition for cell off (0 mV) and cell on (1000 mV) for two different sulfur dioxide doses (0 and 8 g.hL<sup>-1</sup> SO<sub>2</sub>) and histograms illustrating the number of unique detected formulas divided into chemical families, in cell on and cell off, and for 0 and 8 g.hL<sup>-1</sup> SO<sub>2</sub> wine samples.

**Figure 3:** Standardized dataset from electro-oxidation experiments on wines with no SO<sub>2</sub> added, using the Off-line system coupled to FT-ICR-MS, and showing elemental formulas associated with significantly higher intensity according to the voltage applied (0 mV versus 1000 mV). High to low intensity profiles, van Krevelen diagrams (H/C versus O/C) and corresponding sum formula histograms for cell off markers (A) and cell on markers (B). These data were obtained from the hierarchical cluster analysis (ANOVA test, p-values < 0.01). (C) Diagram of the double bond equivalent normalized to the total number of carbon atoms (DBE/C) as a function of the mass range ( $m/z$  100 to 800 Da) with median values highlighted for cell on and cell off markers (D) Diagram of the number of oxygen normalized to the number of hydrogen (O/H) as a function of the mass range ( $m/z$  100 to 800 Da) with median values highlighted for cell on and cell off markers. Elemental formulas in van Krevelen diagrams are colored according to their composition and are sized according to their intensity, CHO in blue, CHOS in green, CHON in orange and CHONS in red.

Oxidation fingerprints obtained from electrochemical simulation coupled to FT-ICR-MS

S- and N-containing compounds were mostly impacted by electrochemical simulation.

Ultrahigh resolution mass spectrometry revealed the corresponding oxidation fingerprints.

This approach paves the way for exploration of potential antioxidation biomarkers.

ACCEPTED MANUSCRIPT

